

Lecture 11

Last time: finished proof of Larson-Radford

§5. character theory and the class equation

$H =$ finite dim'l semisimple Hopf algebra / \mathbb{k} , $\text{char } \mathbb{k} = 0$, $\bar{\mathbb{k}} = \mathbb{k}$.

$\text{Irr}(H) = \{ (V_0, \rho_0), \dots, (V_m, \rho_m) \}$ complete set of ^{representatives} isomorphism classes of irreducible left H -modules. $\rho_i : H \rightarrow \text{End}(V_i)$.

character: $\chi_i := \text{Tr}_{V_i} \circ \rho_i \in H^*$.

Example: We assume $V_0 \cong \mathbb{k} \Lambda$ for $\Lambda \in \int_H$ w/ $\varepsilon(\Lambda) = 1$.

H acts on $V_0 \cong \mathbb{k} \Lambda$ by left multiplication: $\forall h \in H$,

$$L_h(\Lambda) = \underset{\substack{\uparrow \\ \text{multiplication in } H}}{h} \cdot \Lambda = \varepsilon(h) \Lambda, \text{ so } \chi_0(h) = \text{Tr}_{V_0}(L_h) = \varepsilon(h).$$

$$\chi_0 = \varepsilon \in H^*.$$

As noted before, the

$R(H) = \text{span}_{\mathbb{k}} \{ \chi_i \mid 0 \leq i \leq m \} \subseteq H^*$. \vee tensor product of two H -modules

is again an H -module. so for any $0 \leq i, j \leq m$,

$$V_i \otimes V_j \cong \bigoplus_{l=0}^m N_{i,j}^l V_l \quad \text{"fusion rule"}$$

where $N_{i,j}^l = \dim \text{Hom}_H(V_i \otimes V_j, V_l)$ is the multiplicity of V_l in $V_i \otimes V_j$, which are nonnegative integers.

Taking trace on both sides, we have $\chi_i * \chi_j = \sum_{l=0}^m N_{i,j}^l \chi_l$.

This makes $R(H)$ a subalgebra of H^* , called the character algebra.

Note that $1_{R(H)} = 1_{H^*} = \varepsilon$. By Artin-Wedderburn and the uniqueness of trace, $f \in R(H)$ if and only if $f(hk) = f(kh)$ for all $h, k \in H$. (i.e., f is a class function).

For any left H -module V , its linear dual $V^* = \text{Hom}_{\mathbb{K}}(V, \mathbb{K})$ is also a left H -module via $(h \circ f)(v) := f(S(h) \cdot v)$, $\forall v \in V, f \in V^*$.

\uparrow V^* -action \uparrow V -action

By assumption, H is semisimple, $S^2 = \text{id}$, so V^{**} is canonically isomorphic to V .

For any $0 \leq i \leq m$, $V_i^* \in \text{Irr}(H)$, i.e., $\exists 0 \leq i^* \leq m$ s.t. $V_i^* \cong V_{i^*}$.

By definition, $\chi_{i^*} = S^*(\chi_i) = \chi_i \circ S$ for all i .

\uparrow
 antipode of H^*

if $H = \mathbb{K}G$ for a finite group

G , then $\Lambda = \frac{1}{|G|} \sum_{g \in G} g$

$(\varphi | \psi) = \frac{1}{|G|} \sum_{g \in G} \varphi(g) \psi(g^{-1})$

THM 5.1 (Orthogonality of characters)

Choose $\Lambda \in \int_H$ w/ $\varepsilon(\Lambda) = 1$. For any $\varphi, \psi \in H^*$, define

$$(\varphi | \psi) := (\varphi * S^*(\psi))(\Lambda) = \sum \varphi(\Lambda_1) \psi(S(\Lambda_2))$$

then $(\chi_i | \chi_j) = \delta_{i,j}$ for all $0 \leq i, j \leq m$.

Sketch. By the above, $\chi_0 = \varepsilon$, so $\chi_0(\Lambda) = \varepsilon(\Lambda)$. For any $0 < i \leq m$,

$\chi_i(\Lambda) = 0$ since $\Lambda \cdot V_i = 0$. Otherwise, there exists $v \in V_i$ s.t. $\Lambda \cdot v \neq 0$.

Then $\mathbb{K}[\Lambda \cdot v] \subseteq V_i$ is isomorphic to V_0 , which contradicts the choice of V_i .

Now $S^*(\chi_j) = \chi_{j^*}$, so

$$(\chi_i | \chi_j) = (\chi_i * \chi_{j^*})(\Lambda) = \sum_{\ell=0}^m N_{i,j^*}^{\ell} \chi_{\ell}(\Lambda) = N_{i,j^*}^0$$

It is easy to check that the map $\Phi: \text{Hom}_H(V_i \otimes V_{j^*}, V_0) \rightarrow \text{Hom}_H(V_i, V_j)$

determined by $\varepsilon(f(v \otimes w^*)) = w^*(\Phi(f)(v))$ for all $v \in V_i$, $w^* \in V_j^*$ and $f \in \text{Hom}_H(V_i \otimes V_j^*, V_0)$, is a linear isomorphism. By Schur's Lemma,

$$N_{i,j^*}^0 = \dim \text{Hom}_H(V_i \otimes V_j^*, V_0) = \dim \text{Hom}_H(V_i, V_j) = \delta_{i,j}. \quad \square$$

Cor 5.2. For any $0 \leq i, j, l \leq m$, we have

- $(\cdot | \cdot)$ is symmetric and S^* -invariant
- $N_{i,j}^l = (\chi_l | \chi_i \chi_j)$ and
- $N_{i,j}^l = N_{l,i^*j^*} = N_{j,l^*i^*}$

Sketch. The first two statements follow immediately from THM 5.1. For the last one,

$$\begin{aligned} N_{i,j}^l &= (\chi_l | \chi_i \chi_j) = (\chi_l | S^*(\chi_i \chi_j)) = (\chi_l | S^*(\chi_i) S^*(\chi_j)) \\ &= (\chi_l | S^*(\chi_j) S^*(\chi_i)) = (\chi_l \chi_{j^*} | \chi_i) = N_{l,j^*}^i \\ &= (S^*(\chi_l \chi_{j^*}) | S^*(\chi_i)) = (\chi_j \chi_{l^*} | \chi_{i^*}) = N_{j,l^*}^{i^*} \quad \square \end{aligned}$$

Consider $R_{\mathbb{Q}}(H) := \text{span}_{\mathbb{Q}}\{\chi_i \mid 0 \leq i \leq m\}$, then S^* induces a \mathbb{Q} -linear involution

on $R_{\mathbb{Q}}(H)$. We claim that for any $0 \neq \alpha \in R_{\mathbb{Q}}(H)$, $\alpha \alpha^* \neq 0$. Indeed,

write $\alpha = \sum_i a_i \chi_i$ w/ $a_i \in \mathbb{Q}$. then

$$\alpha \alpha^* = \sum_{i,j} a_i a_j \chi_i \chi_{j^*} = \left(\sum_i a_i^2 \right) \varepsilon + \sum_{j>0} b_j \chi_j$$

for some $b_j \in \mathbb{Q}$. Since $\alpha \neq 0$, $\sum_i a_i^2 > 0$, so $\alpha \alpha^* \neq 0$. Suppose the Jacobson radical $J \subseteq R_{\mathbb{Q}}(H)$ is nonzero, then choose $0 \neq \alpha \in J$, and consider $\gamma = \alpha \alpha^* \in J$.

($\gamma \neq 0$ by above), which is nilpotent. Since $\gamma^* = \gamma$, so the above computation

implies $\gamma^2 \neq 0$, and by induction, $\gamma^t \neq 0$ for all $t \geq 1$, contradicting the nilpotency of γ . Thus, $J=0$, which means

LEM 5.3 $R(H)$ is a semisimple algebra. □

Now we can prove the class equation for semisimple finite dim'l Hopf algebras, which is proved for Hopf C^* -algebras by G.I. Kac, and in general by Y. Zha.

THM 5.4. Let H be a finite-dimensional semisimple Hopf algebra over an algebraically closed field \mathbb{k} of characteristic 0. Let $\{e_0, \dots, e_m\}$ be a complete set of primitive orthogonal idempotents in $R(H)$, where e_0 is an integral in H^* , then for each $0 \leq i \leq m$, $\dim(e_i H^*)$ divides $\dim(H)$, and

$$\dim(H) = 1 + \sum_{i=1}^m \dim(e_i H^*).$$

To prove the theorem, we recall the elementary lemma.

symmetric $(x, y) = (y, x)$

LEM 5.5 Suppose $\text{char}(\mathbb{k}) = 0$, $(A, (\cdot, \cdot))$ is a Frobenius algebra / \mathbb{k} of dimension n w/ a dual basis $\{a_j, b_j \mid j=1, \dots, n\}$. Assume $A = \bigoplus_{i=1}^N A_i$

w/ $A_i = M_{n_i}(\mathbb{k})$, and set $f_i := \sum_{j=1}^n \text{Tr}_i(a_j) b_j$, where Tr_i is the

matrix trace in A_i . For any i , if e_i is the primitive central idempotent corresponding to A_i , then $f_i = \alpha_i \cdot e_i$ for some $\alpha_i \in \mathbb{k}^\times$.

Sketch. By assumption, for any $x, y \in A$, $(x, y) = \sum_{t=1}^N \gamma_t \text{Tr}_t(xy)$ for some

$\gamma_t \in \mathbb{k}^x$. Hence, the set $\{ \gamma_t^{-1} u_{r,s}^t, u_{s,r}^t \mid 1 \leq t \leq N, 1 \leq r, s \leq n_t \}$ is a dual basis w.r.t. (\cdot, \cdot) , where $u_{r,s}^t$ is the usual matrix unit in A_t . Moreover, by the properties of dual bases, we have

$$\sum_{t,r,s} \gamma_t^{-1} u_{r,s}^t \otimes u_{s,r}^t = \sum_j a_j \otimes b_j.$$

Therefore,
$$f_i = \sum_{t,r,s} \text{Tr}_i(\gamma_t^{-1} u_{r,s}^t) u_{s,r}^t$$

$$= \sum_r \gamma_i^{-1} \text{Tr}_i(u_{r,r}^i) u_{r,r}^i = \gamma_i^{-1} \sum_r u_{r,r}^i = \gamma_i^{-1} e_i. \quad \square$$

Now we introduce the following twist of the bilinear form (1) on $R(H)$.

Choose $\Lambda \in \int_H$ s.t. $\varepsilon(\Lambda) = 1$. For $\varphi, \psi \in R(H)$, define

$$\langle \varphi, \psi \rangle := (\varphi \psi)(\Lambda) = (\varphi | S^*(\psi))$$

Then $\langle \cdot, \cdot \rangle$ is nondegenerate, associative, and symmetric.

(follows from associativity of \ast)

$$\begin{aligned} & (\varphi | \psi \gamma) \\ &= (\varphi S^*(\psi) | \gamma) \\ & \quad \uparrow \\ & (\varphi \psi | \gamma) \\ & \quad \uparrow \\ & (x_i | x_j x_l) \text{ w/ nec.} \\ &= (x_i x_j \ast | x_l) \text{ assoc.} \\ & (x_i x_j | x_l) \end{aligned}$$

Symmetry follows from Prop 4.11 (2) : since $(S^*)^2 = \text{id}_{H^*}$, then Λ is a nonzero multiple of the character of the regular rep of H^* . i.e., $(\varphi \psi)(\Lambda) = (\psi \varphi)(\Lambda) \forall \varphi, \psi \in H^*$.

Moreover, it follows from THM 5.1 that $\{ S^*(x_j), x_j \}$ is a dual basis for $\langle \cdot, \cdot \rangle$. In fact, $(\langle x_0, \cdot \rangle, S^*(x_j), x_j)$ is a Frobenius system for $R(H)$. (w/ trivial Nakayama).

We present the following proof of THM 5.4 due to Lorenz.

Sketch of PF of THM 5.4. Let $A = R(H) = \bigoplus_{i=1}^N M_{n_i}(K)$, and fix a primitive

idempotent $e = e_i$ in A . We prove the divisibility first. Set

$$d := \frac{\dim(H)}{\dim(eH^*)} \in \mathbb{Q} \subset K.$$

we claim that d is an algebraic integer. (\exists monic polynomial $p(X) \in \mathbb{Z}[X]$

$$\text{s.t. } p(d) = 0). \quad \left. \begin{array}{l} d = \frac{p}{q} \Rightarrow qd - p = 0 \\ \text{if } d \text{ is alg. int, } \text{monic} \end{array} \right\} \Rightarrow |q|=1, d \in \mathbb{Z}.$$

Let E be the central idempotent of A which e belongs to. Then $ER(H) \cong M_{n_i}(K) \cong (eR(H))^{\oplus n_i}$. This implies $EH^* \cong (eH^*)^{\oplus n_i}$. Consequently,

$$\dim(EH^*) = n_i \dim(eH^*), \text{ and so}$$

$$d = \frac{n_i \dim(H)}{\dim(EH^*)}$$

Now let $f = \sum_j \text{Tr}_i(S^*(\alpha_j)) \alpha_j$, then by LEM 5.5, $f = \alpha E$ for some $\alpha \neq 0$

and we claim that $d = \alpha$. Indeed, let $\Lambda \in \int_H$ s.t. $\varepsilon(\Lambda) = 1$, then by Prop 4.11 (a), $\text{Tr}_{H^*} = \dim(H) \cdot \Lambda$. Thus, on the one hand,

\uparrow
 character of regular rep of H^*

$$\dim(EH^*) = \text{Tr}_{H^*}(LE) = \dim(H) \cdot E(\Lambda) = \dim(H) \cdot \alpha^{-1} \cdot f(\Lambda).$$

on the other hand, by THM 5.1,

$$f(\Lambda) = \sum_j \text{Tr}_i(\alpha_j^*) \alpha_j(\Lambda) = \text{Tr}_i(\alpha_0) \alpha_0(\Lambda) = \text{Tr}_i(\varepsilon) \cdot \varepsilon(\Lambda)$$

$$= \text{Tr}_i(\varepsilon) = \eta_i$$

$\hookrightarrow A = \begin{bmatrix} \square & & & \\ & \square & & \\ & & \square & \\ & & & \ddots \end{bmatrix}$, corresponding trace when we view $\varepsilon \in A$.

$$\text{i.e., } \begin{array}{c} \varepsilon \in R(H) \\ \parallel \\ \downarrow_{R(H)} \end{array} \longleftrightarrow \text{id matrix} \in A.$$

Therefore, $d = \frac{\eta_i \dim(H)}{\dim(H) \cdot \alpha^{-1} \cdot \eta_i} = \alpha.$

Now we show that f is integral over \mathbb{Z} . i.e., there is a monic polynomial $p(X) \in \mathbb{Z}[X]$ s.t. $p(f) = 0$. Denote the elements in \mathbb{K} that are integral over \mathbb{Z} by \mathcal{O} , they form a ring called the ring of (algebraic) integers. Consider $R_{\mathbb{Z}}(H) = \mathbb{Z}[\alpha_j \mid 0 \leq j \leq m] \subseteq R(H)$. By orthogonality, $R_{\mathbb{Z}}(H)$ is a finitely generated free \mathbb{Z} -module, so any element $x \in R_{\mathbb{Z}}(H)$ is integral over \mathbb{Z} (see, for example, Neukirch, I. 2).

So all the eigenvalues of x are in \mathcal{O} , which means $\text{Tr}_i(x) \in \mathcal{O}$.

$$R_{\mathbb{Z}}(H) \subset R(H) = A = \bigoplus M_{\eta_j}(\mathbb{K})$$

Thus, all eigenvalues of $f = \sum_j \text{Tr}_i(\alpha_j^*) \alpha_j$ are in \mathcal{O} . So f is integral

over \mathbb{Z} . Let $p(X) \in \mathbb{Z}[X]$ be the monic polynomial s.t. $p(f) = 0$, then

by $f = dE$ and $E^2 = E$, we have $p(d) = 0 \Rightarrow d \in \mathcal{O} \cap \mathbb{Q} = \mathbb{Z}$.

The equality in the theorem follows from $1_{R(H)} = 1_{H^*} = \epsilon = \sum_{j=1}^m e_j$ \square